

CRYSTAL STRUCTURES OF LARGE-VOLUME COMMERCIAL PHARMACEUTICALS

James A. Kaduk, kaduk@polycrystallography.com
North Central College, Naperville IL
Illinois Institute of Technology, Chicago IL
Poly Crystallography Inc., Naperville IL

Austin M. Wheatley
North Central College, Naperville IL

Amy M. Gindhart and Thomas N. Blanton
ICDD, Newtown Square PA

As part of a continuing project, the room-temperature crystal structures of several commercial pharmaceutical APIs have been solved using synchrotron X-ray powder diffraction data (11-BM at APS), and optimized using density functional techniques. All of the structures presented have features which made their solution and refinement non-routine. The molecules to be discussed include: 1. terazosin hydrochloride dihydrate (Hytrin), which was originally solved (but only approximately) using data contained in the Powder Diffraction File database. 2. bretylium tosylate (Bretylol and others), which exhibited significant decomposition in the beam. 3. oxybutynin hydrochloride hemihydrate (Dytropan and Lyrinel XL), which has not been described as a hemihydrate, and which exhibits X-ray induced photoreduction of a triple bond. 4. levocetirizine dihydrochloride (Xyzal), which solves and refines better in $P2_1/n$ rather than the true space group $P2_1$. The presentation may also include progress (or lack thereof) on februxostat Form G (Uloric and Adenuric), as well as other new structures as they are solved.